

Effective Chiral Meson-Baryon Lagrangian from Path Integral Hadronization of a Quark-Diquark Model

D. Ebert* and Th. Jurke†

Institut für Physik, Humboldt-Universität zu Berlin, Invalidenstr. 110, 10115 Berlin, Germany

1st February 2008

Abstract

The approach of path integral hadronization, applied to a quark-diquark toy model, is used to derive an effective chiral meson-baryon Lagrangian. In this approach the Goldberger-Treiman relation, found at the quark-meson level, is re-established at the composite hadron level. For illustrations, electromagnetic low energy characteristics of composite nucleons are estimated.

*e-mail: debert@qft2.physik.hu-berlin.de

†e-mail: jurke@pha2.physik.hu-berlin.de

1 Introduction

It is well-known that low energy properties of mesons and baryons can be suitably described by phenomenological effective Lagrangians [1, 2, 3] which embody the global (chiral) flavour symmetries of quantum chromodynamics ($SU(2) \times SU(2)$ or $SU(3) \times SU(3)$). The appearance of an approximate chiral symmetry and its dynamical breakdown ensures both the generation of Goldstone pions as well as the existence of current algebra relations and low energy theorems.

Concerning meson physics, the approach of path integral bosonization applied to QCD-motivated Nambu - Jona-Lasinio (NJL)-type models [4] turned out to be a powerful method for deriving effective chiral meson Lagrangians [3, 5] (for further references see the recent review [6]).

The treatment of baryons as relativistic bound systems leads us in the case $N_C = 3$ to the concept of diquarks which can form together with a quark a colourless baryon. By applying path integral methods it was in particular possible to derive Faddeev-type equations determining the spectrum of composite baryons [7, 8, 9].

It is further a challenge to employ path integral techniques in collective fields in order to derive effective chiral meson-baryon Lagrangians, including the hadronic Goldberger-Treiman relation, directly from an underlying microscopic quark-diquark interaction. We choose in this letter a simple toy model, containing a local interaction of quarks with (scalar) diquarks in order to demonstrate the powerfulness of such a method of “path integral hadronization”.

Finally, this model is used to get some rough insight into the electromagnetic properties of a composite nucleon.

2 Effective Chiral Meson Lagrangian and Field Transformations

In order to fix our notations required for the derivation of the non-linear meson-baryon Lagrangian it is useful to first recapitulate some results of the

bosonization of the NJL-model [5, 6], emphasizing non-linear field transformations [3]. Thus, we consider the NJL Lagrangian with scalar and pseudoscalar couplings possessing a global flavour and colour $SU(2)_A \times SU(2)_V \times SU(3)_C$ symmetry

$$\mathcal{L}_{\text{NJL}} = \bar{q} \left(i\hat{\partial} - m_0 \right) q + \frac{G}{2} \left[(\bar{q}q)^2 + (\bar{q}i\gamma_5 \vec{\tau}q)^2 \right], \quad (1)$$

where $\hat{\partial} = \partial^\mu \gamma_\mu$, G is a universal coupling constant of dimension $(\text{mass})^{-2}$, $\vec{\tau}$ are the Pauli matrices of the flavour group $SU(2)$ and m_0 is an explicit chiral symmetry-breaking quark mass (summation over repeated indices is always understood). Introducing collective meson fields σ , π_i ($i = 1, 2, 3$) by the Gauß trick

$$e^{i \int d^4x \frac{G}{2} [(\bar{q}q)^2 + (\bar{q}i\gamma_5 \vec{\tau}q)^2]} \equiv \mathcal{N}_1 \int \mathcal{D}\sigma \mathcal{D}\vec{\pi} \times e^{i \int d^4x \left(-\frac{1}{2G} (\sigma^2 + \vec{\pi}^2) - \bar{q}(\sigma + i\gamma_5 \vec{\tau}\vec{\pi})q \right)},$$

one gets a semi-bosonized Lagrangian

$$\mathcal{L}_{\text{NJL}}^{qM} = \bar{q}(i\hat{\partial} - \sigma - i\gamma_5 \vec{\tau}\vec{\pi})q - \frac{1}{2G} \left((\sigma - m_0)^2 + \vec{\pi}^2 \right), \quad (2)$$

where the bare quark mass m_0 has been absorbed into the field σ .

It is convenient to define new scalar and pseudoscalar fields σ' and $\vec{\Phi}$ via the exponential parametrization

$$\sigma + i\gamma_5 \vec{\tau}\vec{\pi} = (m + \sigma') e^{-\frac{i}{F_\pi} \gamma_5 \vec{\tau}\vec{\Phi}},$$

(F_π is the pion decay constant and the constituent quark mass $m \equiv \langle \sigma \rangle_0$ is fixed by the gap equation [5]).

Performing an appropriate chiral rotation of quark fields,¹

¹Wess-Zumino anomalies arising due to this transformation are discarded.

$$q = e^{\frac{i}{F_\pi} \gamma_5 \frac{\vec{\tau}}{2} \vec{\Phi}} \chi,$$

provides us a Lagrange density in terms of redefined fields:

$$\begin{aligned} \mathcal{L}_{\text{NJL}}^{qM}(\chi, \vec{\Phi}, \sigma') = & -\frac{1}{2G}(m + \sigma')^2 \\ & + \frac{m + \sigma'}{16G} m_0 \text{Tr}_{F,D} \left(e^{-\frac{i}{F_\pi} \gamma_5 \vec{\tau} \vec{\Phi}} + \text{h.c.} \right) \\ & + \chi \left\{ i\gamma_\mu \left(\partial^\mu + e^{-\frac{i}{F_\pi} \gamma_5 \frac{\vec{\tau}}{2} \vec{\Phi}} \partial^\mu e^{\frac{i}{F_\pi} \gamma_5 \frac{\vec{\tau}}{2} \vec{\Phi}} \right) \right. \\ & \left. - m - \sigma' \right\} \chi, \end{aligned} \quad (3)$$

where the trace is taken over flavour and Dirac indices.

Note the non-linear transformation law of the meson field $\vec{\xi} \equiv \frac{\vec{\Phi}}{F_\pi}$ under global chiral transformations $g = e^{\frac{i}{2} \gamma_5 \vec{a} \cdot \vec{\tau}} e^{\frac{i}{2} \vec{v} \cdot \vec{\tau}} \in SU(2)_A \times SU(2)_V$ [2, 3] (a_i, v_i are real parameters)

$$g \cdot e^{\frac{i}{2} \gamma_5 \vec{\tau} \vec{\xi}(x)} = e^{\frac{i}{2} \gamma_5 \vec{\tau} \vec{\xi}'(x)} \cdot h(x), \quad (4)$$

where

$$h(x) = e^{\frac{i}{2} \vec{\tau} \vec{u}'(\vec{\xi}(x), g)} \in SU(2)_{V, \text{loc}}$$

is an element of the local vector group.

For later use it is convenient to rewrite the Dirac operator of the χ -field in (3) by employing the Cartan decomposition [3]

$$\begin{aligned} e^{-\frac{i}{2} \gamma_5 \vec{\tau} \vec{\xi}} \partial^\mu e^{\frac{i}{2} \gamma_5 \vec{\tau} \vec{\xi}} = & \\ & \frac{i}{2} \gamma_5 \vec{\tau} \vec{\mathcal{A}}^\mu(\xi) + \frac{i}{2} \vec{\tau} \vec{\mathcal{V}}^\mu(\xi). \end{aligned} \quad (5)$$

The fields $\chi, \vec{\mathcal{A}}_\mu, \vec{\mathcal{V}}_\mu$ have the following simple transformation law under (4) ($\mathcal{V}_\mu \equiv \frac{\vec{\tau}}{2} \vec{\mathcal{V}}_\mu$ etc.)

$$\begin{aligned} \chi & \mapsto \chi' = h(x) \chi \\ \mathcal{V}_\mu & \mapsto \mathcal{V}'_\mu = h(x) \mathcal{V}_\mu h^\dagger(x) - h(x) i \partial_\mu h^\dagger(x) \\ \mathcal{A}_\mu & \mapsto \mathcal{A}'_\mu = h(x) \mathcal{A}_\mu h^\dagger(x). \end{aligned}$$

As we see $\mathcal{V}_\mu$ transforms as a gauge field with respect to $SU(2)_{V, \text{loc}}$. This allows one to define the following chiral-covariant derivative of the rotated quark field χ

$$D_\mu \chi = (\partial_\mu + i \mathcal{V}_\mu) \chi. \quad (6)$$

Thus, using (3), (5) and (6), the inverse propagator of the χ -field takes the form

$$S^{-1} = i \hat{D} - m - \sigma' - \hat{\mathcal{A}} \gamma_5. \quad (7)$$

Finally, by integrating over the χ -field in the generating functional of (3) and freezing the σ' -field, one arrives at a non-linear pion Lagrangian. Indeed, performing the loop expansion of the resulting quark determinant ($\det S^{-1}$) and choosing a gauge-invariant regularisation, we arrive at a mass-like term of the $\mathcal{A}_\mu(\xi)$ -field contributing to $\mathcal{L}_{\text{eff}}^M$,²

$$\mathcal{L}_{\text{eff}}^M = \frac{m^2}{g_{\pi qq}^2} \text{Tr}_F \mathcal{A}_\mu^2 + \Delta \mathcal{L}_{\text{sb}}.$$

Here $g_{\pi qq}$ is the induced meson-quark coupling constant and $\Delta \mathcal{L}_{\text{sb}} = \mathcal{O}(m_0)$ is the symmetry-breaking mass term given by the second term in (3). As discussed in [3] the Cartan form $\mathcal{A}_\mu(\xi)$ is just the chiral covariant derivative of the ξ -field, admitting the expansion

$$\begin{aligned} \vec{\mathcal{A}}_\mu(\xi) & \equiv D_\mu \vec{\xi} = \partial_\mu \vec{\xi} + \mathcal{O}(\xi^3) \\ & = \frac{1}{F_\pi} \partial_\mu \vec{\Phi} + \mathcal{O}(\Phi^3). \end{aligned}$$

Thus, using the Goldberger-Treiman relation,

$$F_\pi = \frac{m}{g_{\pi qq}},$$

at the quark level [5], one obtains the effective chiral meson Lagrangian

$$\mathcal{L}_{\text{eff}}^M = \frac{F_\pi^2}{2} D_\mu \vec{\xi} D^\mu \vec{\xi} + \Delta \mathcal{L}_{\text{sb}}. \quad (8)$$

²Note that possibly arising field strength terms of the $\mathcal{V}_\mu$ - and $\mathcal{A}_\mu$ -fields are vanishing.

3 Hadronization of a Quark-Diquark Toy Model

In order to derive an effective chiral meson-baryon Lagrangian from a microscopic quark-diquark model, the semi-bosonized Lagrangian (3) has to be supplemented by the kinetic part of the scalar isoscalar diquark D , which here will be treated as an elementary field, and by a quark-diquark interaction term. Let us consider a simple toy model defined by the following extended chiral invariant Lagrangian,

$$\mathcal{L}^{qMD} = \mathcal{L}_{\text{NJL}}^{qM} + D^\dagger \Delta^{-1} D + \tilde{G} \bar{\chi} D^\dagger D \chi \quad (9)$$

$$\Delta^{-1} = -\partial_\mu \partial^\mu - M_D^2, \quad (10)$$

with diquark mass M_D and a local quark-diquark interaction with coupling constant $\tilde{G}$.

As in the NJL model we are now in the position to introduce collective baryon fields B using a Gauß trick. Here we have

$$e^{i \int d^4x \tilde{G} \bar{\chi} D^\dagger D \chi} = \mathcal{N}' \int \mathcal{D}B \mathcal{D}\bar{B} e^{i \int d^4x \left(-\frac{1}{\tilde{G}} \bar{B} B - \bar{\chi} D^\dagger B - \bar{B} D \chi \right)}. \quad (11)$$

Let us next perform the “hadronization” of the Lagrangian (9) by integrating step by step over quark and diquark fields in the respective generating functional:

$$Z = \mathcal{N}_1 \int \mathcal{D}\sigma' \mathcal{D}\Phi_i \mathcal{D}B \mathcal{D}\bar{B} D D D D^\dagger \times e^{i \int d^4x \left[-i \text{Tr}_{F,D} \ln S^{-1} - \frac{1}{\tilde{G}} \bar{B} B \right]} \times e^{i \iint d^4x d^4y \left[D^\dagger(x) (\Delta^{-1} - \bar{B} S B)_{(x,y)} D(y) \right]},$$

$$Z = \mathcal{N}_2 \int \mathcal{D}\sigma' \mathcal{D}\Phi_i \mathcal{D}B \mathcal{D}\bar{B} \times \exp \left\{ i \int d^4x \left\{ -i \text{Tr}_{F,D} \ln S^{-1} - \frac{1}{\tilde{G}} \bar{B} B + i \ln(1 - \bar{B} S \Delta B)_{(x,x)} \right\} \right\}, \quad (12)$$

with the quark and diquark propagators S and Δ defined in (7) and (10), respectively.

Expanding the logarithms in power series at the one-loop level (see Fig. 1) and taking into account only lowest order derivative terms, one describes both the generation of kinetic and mass terms for the composite baryon fields as well as the meson-baryon interaction. This yields the expression

$$\mathcal{L}_{\text{eff}}^{MB} = \int d^4x \int d^4y \bar{B}(x) \times \left[\left(-\frac{1}{\tilde{G}} - Z_1^{-1} \gamma_\mu \frac{\vec{\tau}}{2} \vec{V}^\mu(x) - g_A \gamma_\mu \gamma_5 \frac{\vec{\tau}}{2} \vec{A}^\mu(x) \right) \delta^4(x-y) - \Sigma(x-y) \right] B(y), \quad (13)$$

where the baryon self-energy Σ admits in momentum space the decomposition $\Sigma(p) = \hat{p} \Sigma_V(p^2) + \Sigma_S(p^2)$, whose low-momentum expansion will be quoted below; Z_1^{-1} and g_A are the vector vertex renormalization and axial coupling constant defined by suitably regularized loop integrals.




Fig. 1: Self-energy contribution of the nucleon, vector vertex diagram and axial-vector vertex diagram resulting from expanding the second logarithm in Eq. (12) of the text.

Performing the low-momentum expansion of Σ and introducing renormalized fields we have

$$\mathcal{L}_{\text{eff}}^{MB} = \bar{B}_{\text{ren}} \left(i \hat{D} - M_B \right) B_{\text{ren}} - g_A^{\text{ren}} \bar{B}_{\text{ren}} \gamma_\mu \gamma_5 \frac{\vec{\tau}}{2} B_{\text{ren}} \vec{A}^\mu, \quad (14)$$

with $B = Z^{\frac{1}{2}} B_{\text{ren}}$, $g_A^{\text{ren}} = Z g_A$ and Z being the

Z -factor derived from the baryon propagator which satisfies the Ward identity $Z = Z_1$.

As seen from (13) the nucleon mass is estimated by

$$\frac{1}{\bar{G}} + \left(M_B \Sigma_V(M_B^2) + \Sigma_S(M_B^2) \right) = 0.$$

Recalling $\mathcal{A}_i^\mu = \partial^\mu \xi_i + \dots$, we obtained an axial-vector derivative coupling of pions to baryons. – To get rid of the derivative let us redefine the baryon fields via

$$B_{\text{ren}} = e^{-\frac{i}{2} g_A^{\text{ren}} \gamma_5 \vec{\tau} \vec{\xi}} \tilde{B}.$$

Then, the expansion in a power series in $\vec{\Phi}$ provides us the expression

$$\begin{aligned} \mathcal{L}_{\text{eff}}^{MB} &= \bar{\tilde{B}} \left(i \hat{\partial} - M_B \right) \tilde{B} \\ &+ g_A^{\text{ren}} \frac{M_B}{F_\pi} \bar{\tilde{B}} i \gamma_5 \vec{\tau} \tilde{B} \vec{\Phi} + \mathcal{O}(\Phi^2). \end{aligned} \quad (15)$$

Obviously we have to identify the constant, associated to the Yukawa interaction, with the pion-nucleon coupling constant as

$$g_{\tilde{B}\tilde{B}\Phi}^{\text{ren}} = \frac{M_B}{F_\pi} g_A^{\text{ren}}, \quad (16)$$

which is nothing but the Goldberger-Treiman relation at the composite hadron level. Combining (8) and (15), the total meson-baryon Lagrangian reads

$$\mathcal{L}_{\text{eff,tot}}^{MB} = \mathcal{L}_{\text{eff}}^M + \mathcal{L}_{\text{eff}}^{MB}.$$

4 Numerical Discussion

As a simple application let us calculate the renormalized axial coupling constant from the matrix element associated to the third Feynman diagram of Fig. 1,

$$\begin{aligned} iM_\mu^{g_A} &= \\ &- 3i \int iS(p-k) \gamma_\mu \gamma_5 iS(p-k) i\Delta(k) \frac{d^4 k}{(2\pi)^4}. \end{aligned}$$

For typical quark and diquark constituent masses (we suppose an exact isospin symmetry $m_u = m_d = m$) used in such type of approaches

$$\begin{aligned} m &= 0.450 \text{ GeV} \\ M_D &= 0.650 \text{ GeV} \end{aligned}$$

and a cutoff

$$\Lambda = 0.750 \text{ GeV}$$

the model leads to a value

$$g_A^{\text{ren}} = 1.07$$

which is in accord with other related studies (see *e.g.* [10]) but lower then the experimental value $g_A = 1.26$.

By introducing electromagnetic interactions into the model it is straightforward to evaluate electromagnetic nucleon form factors (see Fig. 2).




Fig. 2: Vertex diagrams contributing to the electromagnetic form factors of the nucleon.

With the above mentioned parameter set and by taking a diquark form factor ($q = p' - p$),

$$F_D(q^2) = (1 + 1.05 \text{ GeV}^{-2} \cdot q^2),$$

suggested in [11], our toy model with scalar diquarks leads to the following estimates for proton and neutron magnetic moments as well as electric and magnetic radii, respectively [12]:

	μ_B	exp.	$\langle r^2 \rangle_e$ [fm ²]	exp. [fm ²]	$\langle r^2 \rangle_m$ [fm ²]	exp. [fm ²]
P	1.66	2.79	0.29	0.74	0.20	0.74
N	-0.76	-1.91	-0.01	-0.12	0.19	0.77

Table 1: Estimates within the quark-diquark model for proton and neutron magnetic moments as well as electric and magnetic radii.

The numerical considerations of the relevant Feynman diagrams yield only a weak dependence on constituent masses and on the special choice of the cutoff.

As a further illustration we want to present here the low energy dependence of Sachs' electric form factors of the proton and neutron, respectively (experimental data are taken from [13]).




Fig. 3: Proton electric form factor.




Fig. 4: Neutron electric form factor.

Concerning the estimates shown in the Tab. 1 and Fig. 3, 4, it is expected that the observed discrepancies in the predictions of the toy model from experimental data disappear, if one uses a more realistic, but also more complicated model including both non-local quark-diquark interactions due to quark exchange [7, 8, 9] and axial-vector diquarks.

5 Summary

By applying the path integral hadronization method to a simple local quark-diquark toy model (with scalar diquarks) we have derived an effective chiral meson-baryon Lagrangian. As an important result the Goldberger-Treiman relation at the composite meson-baryon level has been re-established.

For illustrations, we have further given some numerical results concerning the axial coupling constant and low-energetic electric and magnetic form factors of nucleons. A better quantitative correspondence with data obviously demands the use of non-local quark-diquark interactions as well as the additional inclusion of the spin carrying axial diquark. Especially the large differences in the magnetic properties (while getting relations $\mu_{\text{prot}}/\mu_{\text{neutr}}$ close to experi-

mental findings) are strong hints in this direction; a statement which is in accord with other recent papers about this topic [14]. – This could be a very promising subject of further investigations.

Acknowledgements

We wish to thank Th. Feldmann, G. Hellstern and V. Keiner for useful discussions.

References

- [1] S. Weinberg, Phys. Rev. Lett. **18**, 188 (1967).
- [2] S. Coleman, J. Wess and B. Zumino, Phys. Rev. **177**, 2239 (1969);
C. Callan, S. Coleman, J. Wess and B. Zumino, Phys. Rev. **177**, 2247 (1969).
- [3] D. Ebert and M.K. Volkov, Fortschr. Phys. **29**, 35 (1981).
- [4] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961);
Y. Nambu and G. Jona-Lasinio, Phys. Rev. **124**, 246 (1961).
- [5] D. Ebert and M.K. Volkov, Z. Phys. **C16**, 205 (1983);
D. Ebert and H. Reinhardt, Nucl. Phys. **B271**, 188 (1986).
- [6] D. Ebert, H. Reinhardt and M.K. Volkov, Prog. Part. Nucl. Phys. **33**, 1 (1994).
- [7] R.T. Cahill, C.D. Roberts and J. Praschifka, Aust. J. Phys. **42**, 129 (1989).
- [8] H. Reinhardt, Phys. Lett. **B244**, 316 (1990).
- [9] D. Ebert and L. Kaschluhn, Phys. Lett. **B297**, 367 (1992).
- [10] G. Hellstern and C. Weiss, Phys. Lett. **B351**, 64 (1995).
- [11] C. Weiss, A. Buck, R. Alkofer and H. Reinhardt, Phys. Lett. **B312**, 6 (1993).
- [12] Th. Jurke, Diploma thesis, Humboldt-Universität zu Berlin, 1997.
- [13] T. Eden *et al.*, Phys. Rev. **C50**, R1749 (1994);
G. Höhler *et al.*, Nucl. Phys. **B114**, 505 (1976);
M. Meyerhoff *et al.*, Phys. Lett. **B327**, 201 (1994);
S. Platchkov *et al.*, Nucl. Phys. **A510**, 740 (1990).
- [14] V. Keiner, Z. Phys. **A354**, 87 (1996).